

# DEATH STATISTICS

## CRUDE DEATH RATE

In 2000, the crude death rate for Lancaster County was 6.6 per 1,000 population. The age adjusted death rate for this year was 7.65 per 1,000 population. The average crude death rate between 1981 and 2000 was 6.98 per 1,000 population.

The crude death rate and age adjusted death rate for the State of Nebraska during 2000 was 8.8 and 7.9 per 1,000 population respectively. Figure 14 illustrates the death rates for Lancaster County and the State of Nebraska.

## TOTAL NUMBER OF DEATHS

There were 1,653 deaths among Lancaster County residents during 2000. Of these, 1,528 (92.4 %) were residents of Lincoln. The remaining 125 (7.5 %) resided outside the City of Lincoln. The most frequent

months for death were January followed by November (Table 14).

### Race

Of the deaths in 2000, 1,607 (97.2 %) were among Whites, 24 were among Blacks, 17 were among American Indians, and 12 were among Asians. Non-White deaths accounted for 2.7 percent of total deaths. Crude death rate was 7.12 per 1,000 for whites, 3.4 per 1,000 for Blacks, 5 per 1,000 for American Indians and 1.67 per 1,000 for Asians.


### Sex

During 2000, 47.1 percent of all deaths in Lancaster County were males, and 52.9 percent were females.

### Age

In 2000, 22 infants (less than 1 year of age) died, accounting for 1.33 percent of all deaths and an infant mortality rate of 5.9 per

**Figure 14: Comparison of Death Rates\***  
Lancaster County and Nebraska  
1981 to 2000



\*Rate per 1000 Population  
Lincoln-Lancaster Health Department

1,000 live births (*See page 31 for further discussion of infant mortality*). A total of 31 children in the 0-14 year age group died in 2000, accounting for 1.9 percent of all deaths. The most frequent age group for deaths was 80 to 89 years (30.4 % of all deaths) followed by 70 to 79 years (24.3 % of all deaths) (Table 16).

In 2000, the mean age of death for males and females was 69.7 and 77.8 years, respectively (Figure 15). For males, this represents an increase of 2.1 years over the past decade (since 1990). The mean age of death for females increased 1.5 years from 1990 to 2000.

Using a three year rolling average death rate (1998 to 2000), life expectancy at birth (average number of years that an infant would live) in the Lancaster County was 79.1 years (Table 28). Female (81.4 years)

had a higher life expectancy than Male (76.7 years). The overall Life expectancy was 79.1 years for Whites, 77.3 years for Blacks, 64 years for American Indians and 85.2 years for Asians.

## LEADING CAUSES OF DEATH

Except for 1999, heart disease has always been the leading cause of death in Lancaster County . In 2000, a total of 416 deaths were due to Heart Disease and 408 were due to Cancer. These two major causes of death accounted for half of all deaths (Figure 16). Table 17 details statistics on the leading causes of death in Lancaster County.

Table 15 lists the average age of death by sex by selected cause. The six leading causes of death in Lancaster County and Nebraska can be found in Table 18.

**TABLE 14**  
**Deaths by Month**  
**Lancaster County 2000**

Month	No. Deaths	Percent
January	182	11
February	126	7.6
March	160	9.7
April	130	7.9
May	137	8.3
June	100	6
July	127	7.7
August	127	7.7
September	106	6.4
October	152	9.2
November	164	9.9
December	142	8.6
<b>Total</b>	<b>1653</b>	<b>100</b>

Lincoln-Lancaster County Health Department

**TABLE 15.**  
**Average Age of Death**  
**by Sex by Selected Cause**  
**Lancaster County, 2000.**

Cause	Male	Female
Heart Disease	75.9	82.4
Cancer	70.7	72
Cerebrovascular Disease	80.2	82.6
Accidental Deaths	47.3	66.4
Chronic Lung Disease	76.2	77.2
Diabetes Mellitus	66.6	79
Suicide	39.8	48.7

Lincoln-Lancaster County Health Department

## Leading Causes of Death by Age and Gender and Race

Among males, Cancer was the number one cause of death in 2000 followed by Heart Disease and Accident (Table 19). For females the leading cause of death was Heart Disease followed by Cancer and Cerebrovascular Diseases. The same gender difference for leading causes of death was observed in 1999.


Leading Causes of death among different age and racial groups are presented in Table 27 and 28 respectively. Accident was the predominant cause of death among 5 to 44 years age groups.

Among age group 45 to 74 years, cancer was the leading cause of death whereas the leading cause of death for 75 years and older was heart disease.

Causes of infant mortality are listed in Table 23.

**Figure 15: Mean Age of Death for Males and Females**

Lancaster County, NE  
1981-2000



**TABLE 16.**  
**Deaths by Age and Sex**  
**Lancaster County, 2000**

Age	Total Deaths	Percent	Male	Percent of Male	Female	Percent of Female
<1	22	1.3	10	1.3	12	1.4
1-4	3	0.2	3	0.4	0	0
5-9	1	0.1	0	0	1	0.1
10-14	5	0.3	5	0.6	0	0
15-19	9	0.5	6	0.8	3	0.3
20-24	8	0.5	8	1	0	0
25-29	11	0.7	8	1	3	0.3
30-39	42	2.5	31	4	11	1.3
40-49	79	4.8	49	6.3	30	3.4
50-59	99	6	55	7.1	44	5
60-69	194	11.7	116	14.9	78	8.9
70-79	402	24.3	209	26.8	193	22.1
80-89	503	30.4	207	26.6	296	33.9
90-99	255	15.4	68	8.7	187	21.4
>100	20	1.2	4	0.5	16	1.8
<b>Total</b>	<b>1653</b>	<b>100</b>	<b>779</b>	<b>100</b>	<b>874</b>	<b>100</b>

**Figure 16: Leading Causes of Death**  
Lancaster County, NE  
2000




TABLE 17.  
Leading causes of Death  
and Crude Rate per 100,000 Population  
Lancaster County, 2000

Cause of Death	Number of Total Deaths	Percent of Total Deaths	Crude Death Rate*	Age Adjusted Death Rate*
Heart Disease	416	25.2	166.2	194.5
Cancer	408	24.7	163	192.8
Cerebrovascular Disease	105	6.4	42	49
Accidental Deaths	78	4.7	31.2	32.3
Chronic Lung Disease	76	4.6	30.4	36.3
Diabetes Mellitus	56	3.4	22.4	26.3
Pneumonia	43	2.6	17.2	20.1
Alzheimer's disease	39	2.3	15.6	17.9
Suicide	28	1.7	11.2	11.1
Nephritis and Nephrosis	22	1.3	8.8	10.4
Atherosclerosis	20	1.2	8	9.3
Birth defects	11	0.7	4.4	4.4
Essential Hypertension and Hypertensive Renal Disease	11	0.7	4.4	5
Parkinson's Disease	11	0.7	4.4	5.1
Anaemias	10	0.6	4	4.5
Emphysema	10	0.6	4	4.9
Pneumonitis due to solids and liquids	9	0.5	3.6	4.1
Aortic Aneurysm	7	0.4	2.8	3.4
Chronic Liver Disease and Cirrhosis	7	0.4	2.8	2.9
Septicemia	7	0.4	2.8	3.3
Peptic Ulcer	6	0.4	2.4	2.8
Asthma	4	0.2	1.6	1.9
HIV	4	0.2	1.6	1.6
Cholelithiasis and other Disorders of the Gall Bladder	3	0.2	1.2	1.4
Nutritional deficiencies	3	0.2	1.2	1.3
Assult (Homicide)	3	0.2	1.2	0.1
Hernia	2	0.1	0.8	0.1
<b>Appendicitis</b>	<b>1</b>	<b>0.1</b>	<b>0.4</b>	<b>0.5</b>
Influenza	1	0.1	0.4	0
Tuberculosis	1	0.1	0.4	0.5
Viral hepatitis	1	0.1	0.4	0.3
All other causes	250	15	99.8	114
Total	1653	100	6.6	7.7


Lincoln-Lancaster County 2000

TABLE 18.  
Comparison of the Six Leading Causes of Death  
Lancaster County vs. the State of Nebraska, 2000

Lancaster County		Nebraska	
Cause of Death	Percent	Cause	Percent
Heart Disease	25.2	Heart Disease	27.9
Cancer	24.7	Cancer	22.5
Cerebrovascular Disease	6.4	Cerebrovascular Diseases	7.3
Accidental Deaths	4.7	Chronic Lung Disease	4.8
Chronic Lung Disease	4.6	Accidental Deaths	4.3
Diabetes Mellitus	3.4	Diabetes Mellitus	2.7

Lincoln-Lancaster County Health Department

Figure 17: Infant Mortality Rate  
Lancaster County, NE  
1984-2000



Lincoln-Lancaster County Health Department

## Cancer Sites

Cancers of the lung, trachea and bronchus were the leading cause of cancer-related death among males, and females (Table 20). Lymphoid and colon/rectum cancer were the next leading causes of cancer death in men. Lung, trachea, and bronchus cancer were the leading causes of cancer death among women (Table 20). Cancer death rates are discussed on page 34.

## Suicide

Suicide by method and age category are shown in Table 21. The average age for male suicides was 39.8 years, compared to 48.6 years for females. Over three fourths of suicide deaths (80.0 per cent) were among individuals 39 years of age or younger. In 2000, the crude rate for suicide was 11.2 per 100,000 population (Age adjusted rate 11.1 per 100,000). This indicates a remarkable change from the 1999 rate of 6.3 per 100,000 population. The Lancaster County Health Objective for the Year 2010 is to reduce suicide mortality to an age-adjusted rate of 7.5 per 100,000 population.

## INFANT MORTALITY RATE

In 2000, 22 infant deaths occurred in Lancaster County for an infant mortality rate of 5.9 per 1,000 live births (Figure 17). This represents a decrease from the 1999 infant mortality rate (6.4 per 1,000) in which 23 infant deaths occurred. The Lancaster County Health Objective for the Year 2010

is 5.0 per 1,000 live births. In 2000, 45.5% of infant deaths were male while the remaining 54.5% were female. A majority (86.4 per cent) of all infant deaths in 2000 were white (Table 22), a rate of 5.6 per 1,000 white births. Three infant deaths among other races in 2000 translated into an infant mortality rate of 8 per 1,000 non-white births. The leading cause of infant mortality was congenital anomalies (Table 23)

TABLE 19.  
Six Leading Causes of Death by Gender  
Lancaster County 2000

Rank	Gender	
	Male	Female
1	Cancer (214)* 171.1 **	Heart Disease (226) 180.4
2	Heart Disease (190) 151.9	Cancer (194) 155.2
3	Accidental Deaths (47) 37.6	Cerebrovascular Diseases (71) 56.8
4	Chronic Lung Disease (38) 30.4	Chronic Lung Disease (38) 30.4
5	Cerebrovascular Diseases (34) 27.2	Diabetes Mellitus (34) 27.2

\* Number of Deaths    \*\* Sex Specific Rate per 100,000 Population



**TABLE 20.**  
**Cancer by Type**  
**Lancaster County, 2000**

Type of Cancer	ICD 10	Male	Female	Total	Percentage
Lung, Trachea, Bronochus	C33-C34	68	54	122	29.9
Colon/ Rectum	C18-C21	25	24	49	12
Lymphoid and Histiocytic	C81-C96	32	23	55	13.5
Breast	C50	0	23	23	5.6
Prostate	C61	22	0	22	5.4
Pancreas	C25	9	9	18	4.4
Ovary/other Uterine adnexa	C56-C57	0	11	11	2.7
Brain	C71	5	4	9	2.2
Esophagus	C15	10	3	13	3.2
Stomach	C16	7	2	9	2.2
Bladder	C67	7	2	9	2.2
Liver	C22	2	3	5	1.2
Cervix Uteri	C53	0	0	0	0
Skin	C43	2	2	4	1
All other causes	...	25	34	59	14.5
Total	...	214	194	408	100

*Lincoln-Lancaster County Health Department*

**TABLE 21.**  
**Suicides by Type by Age Group**  
**Lancaster County, 2000.**

Types of Suicide	0-14	15-29	30-44	45-59	60 & Over	Total
Fire arms and explosives	1	3	4	2	3	13
Hanging and Suffocation	0	2	3	1	1	7
Other And Unspecified Means	0	1	0	0	0	1
Poisoning by Solids or Liquids	0	0	3	2	0	5
Poisoning by Other Gases and Vapors	0	0	1	0	1	2
<b>Total</b>	<b>1</b>	<b>6</b>	<b>11</b>	<b>5</b>	<b>5</b>	<b>28</b>

*Lincoln-Lancaster County Health Department*

## FETAL DEATHS

Sixteen fetal deaths were reported in 2000, representing a fetal death ratio of 4.3 per 1,000 live births (Figure 18). This is an decrease from the 21 fetal deaths reported during 1999, in which a fetal death ratio of 5.8 per 1,000 live births occurred.

## CRUDE AND AGE-ADJUSTED RATES FOR THE THREE LEADING CAUSES OF DEATH

### Heart Disease

In 2000, 416 (25.2 per cent of deaths) Lancaster County residents died of heart disease. These 416 deaths translate into a crude death rate of 166.2 per 100,000 population. The average crude death rate for heart disease from 1996 through 2000 was 193.7 per 100,000. The Nebraska Department of Health and Human Services reported an average crude death rate for heart disease of 282.8 per 100,000 population for the State of Nebraska between 1996 and 2000.

The 2000 Nebraska Vital Statistics Report lists Nebraska's age-adjusted death rate for heart disease as 216.8 per 100,000 population for 2000, and 242.6 per 100,000 for 1996 through 2000. Using the same age-adjusted method, Lancaster County's age-adjusted death rates for heart disease were lower: 194.5 per 100,000 for 2000 and 212.2 per 100,000 for 1996-2000.

**TABLE 22.**  
Infant Deaths by Race and Sex  
Lancaster County, 2000.

Race	Male	Female	Total
White	8	11	19
Black	1	1	2
American Indian	0	0	0
Asian	1	0	1
Total	10	12	22

*Lincoln-Lancaster County Health Department*

**TABLE 23.**  
Infant Deaths by Cause and Sex  
Lancaster County, 2000

Cause of Death	Male	Female	Total
Congenital Anomalies	4	2	6
Sudden Infant Death Syndrome	2	0	2
Complications during pregnancy	0	3	3
Other	4	7	11
Total	10	12	22

*Lincoln-Lancaster County Health Department*

## Cancer


In 2000, 408 (24.7 % of total deaths) Lancaster County residents died of cancer. These 408 deaths translate into a crude death rate due to cancer of 163 per 100,000 estimated population. This compares to an average crude death rate of 165.9 per 100,000 population for 1996 through 2000, which is lower than the average crude death rate due to cancer of 201.2 per 100,000 estimated population for the State of Nebraska between 1996 and 2000.

The Nebraska Vital Statistics Report lists Nebraska's age-adjusted death rate for cancer as 186.1 per 100,000 population for 2000, and 183.8 per 100,000 for 1996 through 2000. It cites lower age adjusted rates for Lancaster County: 192.8 per 100,000 for 2000 and the same average rate (183.3) of the state per 100,000 for 1996 through 2000.

## Cerebrovascular Disease

In 2000, 105 (6.4 % of total deaths) Lancaster County residents died of cerebrovascular disease. These 105 deaths translate into a crude death rate due to cerebrovascular disease of 42 per 100,000 estimated population. This compares to an average crude death rate of 47.5 per 100,000 estimated population for 1996 through 2000. The average crude death rate due to cerebrovascular disease of 68.3 per 100,000 estimated population for the State much higher than that of Lancaster County. The 2000 Nebraska Vital Statistics Report states that for the State of Nebraska the age-adjusted death rate for cerebrovascular disease was 55.8 per 100,000 population for 2000, and 57.8 per 100,000 for 1996 through 2000. Rates for Lancaster County were lower: 49 per 100,000 for 2000 and 52 per 100,000 for 1996 through 2000.

**Figure 18: Fetal Death Rate**  
Lancaster County, NE  
1981-2000



Lincoln-Lancaster County Health Department

*\*Note: To compare death rates from different populations (Lancaster County and the State of Nebraska), age-adjusted rates are used.*

## DEATHS DUE TO UNINTENTIONAL INJURY

In 2000, 78 deaths occurred (4.7 % of total deaths) due to unintentional injury in Lancaster County. The average ages for male and female deaths were 47.3 and 66.4 years, respectively. Males represented 60.3 percent of all deaths, compared to 39.7 percent for females.

Falls, in recent years for the first time became the leading cause of deaths (28.2 %) due to unintentional injury in 2000, followed by Motor vehicle traffic crashes (26.9 %). Unintentional injury deaths by cause is presented in Table 24.

## DISPOSITION AND PLACE OF DEATH

The most frequent disposition of a body was burial (69.6 %), followed by cremation (27.8 %) (Table 25). The most frequent place of death was in a nursing home, (37.6 %), followed by a hospital or medical center-inpatient (29.7 %) (Table 26).

**TABLE 24.**  
Unintentional Injury Deaths by Principal Cause  
Lancaster County, 2000.

Principal Cause	Frequency
Falls	22
Motor Vehicle Traffic Crash	21
Submersion, Suffocation, or Foreign Bodies	12
Poisoning by Drugs	5
Other Unintentional Injury	18
Total	78

Lincoln-Lancaster County Health Department

**TABLE 25.**  
Disposition of Body  
Lancaster County, 2000

Disposition	Frequency	Percent
Burial	1151	69.6
Cremation	459	27.8
Removal	20	1.2
Donation	20	1.2
Unknown	3	0.2
Total	1653	100

Lincoln-Lancaster County Health Department

**TABLE 26.**  
Place of Death  
Lancaster County 2000.

Place of Death	Frequency	Percent
Nursing Home	622	37.6
Hospital or Medical Center Inpatient	491	29.7
Residence	377	22.8
Hospital or Medical Center Outpatient/ER	97	5.9
Other Facilities	64	3.9
Dead on Arrival at Hospital or Medical Center	1	0.1
Unknown	1	0.1
Total	1653	100

Lincoln-Lancaster County Health Department

**Table 27 Leading Causes of Death By Age  
Lancaster County 2000**

Rank	Age Category										
	Under1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85+
<b>1</b>	Congenital Anomalies 6* (166.6)**	Congenital Anomalies 1 (7.6)	Accident 2 (6.2)	Accident 12 (24.7)	Suicide 7 (18.3)	Accident 12 (31.6)	Cancer 38 (115.9)	Cancer 54 (300.9)	Cancer 112 (840.2)	Heart Disease 134 (1439.3)	Heart Disease 167 (4854.6)
<b>2</b>	Bacterial Sepsis of New Born 3 (83.3)	...		Suicide 4 (8.2)	Accident 6 (15.7)	Cancer 12 (31.6)	Heart Disease 15 (45.7)	Heart Disease 26 (144.9)	Heart Disease 64 (480)	Cancer 128 (1374.8)	Cancer 62 (1802)
<b>3</b>	Complication of Pregnancy 3 (83.3)	...	...	...	Heart Disease 2 (5.2)	Heart Disease 8 (21)	Accident 8 (24.4)	Diabetes Mellitus 8 (44.6)	Chronic Lung Disease 17 (127.5)	Chronic Lung Disease 36 (386.7)	Cerebrovascular Diseases 50 (1453.5)
<b>4</b>	Sudden Infant Death Syndrome 2 (55.5)	...	...	...	Diabetes Mellitus 2 (5.2)	Suicide 6 (15.8)	Suicide 5 (15.2)	Chronic Lung Disease 6 (33.4)	Cerebrovascular Diseases 15 (112.5)	Cerebrovascular Diseases 35 (375.9)	Alzheimer Disease 29 (843)

\* Number of Deaths

\*\* Age Specific Rate per 100,000 Population

**TABLE 28**  
**Five Leading Causes of Death by Race**  
**Lancaster County 2000**

Rank	Race			
	White	Black	American Indian	Asian
1	Heart Disease 409* (181.4)**	Cancer 6(85.1)	Accident 3 (187.6)	Cancer 5 (69.8)
2	Cancer 395 (175.2)	Cerebrovascular Disease 3 (42.5)	Cancer 2 (125.1)	Heart Disease 2 (27.9)
3	Cerebrovascular Disease 102 (45.2)	Diabetes Mellitus 3 (42.5)	Assult 2 (125.1)	...
4	Accident 75 (33.3)	Heart Disease 2 (28.4)	Heart Disease 1 (62.5)	...
5	Chronic Lung Disease 76 (33.7)	...	...	...

\* Number of Deaths    \*\* Race Specific Rate per 100,000 Population  
*Lincoln-Lancaster County Health Department*

**TABLE 29**  
**Life Table For Lancaster Population**

Age groups	Average Remaining Lifetime
<1	79.1
1to4	78.7
5to9	74.7
10to14	69.8
15to19	64.9
20to24	60
25to29	55.1
30to34	50.2
35to39	45.4
40to44	40.7
45to49	36.1
50to54	31.5
55to59	27
60to64	22.6
65to69	18.8
70to74	15.3
75to79	12
80to84	9.2
85+	6.8

**TABLE 30**  
**Life Table For Lancaster Male Population**

Age groups	Average Remaining Lifetime
<1	76.7
1to4	76.2
5to9	72.3
10to14	67.4
15to19	62.5
20to24	57.7
25to29	52.8
30to34	48
35to39	43.3
40to44	38.6
45to49	34
50to54	29.5
55to59	25.1
60to64	20.8
65to69	17
70to74	13.8
75to79	10.7
80to84	8.1
85+	6.2

**TABLE 31**  
**Life Table For Lancaster Female Population**

Age groups	Average Remaining Lifetime
<1	81.4
1to4	80.9
5to9	76.9
10to14	72
15to19	67
20to24	62.1
25to29	57.1
30to34	52.2
35to39	47.4
40to44	42.6
45to49	37.8
50to54	33.2
55to59	28.6
60to64	24.2
65to69	20.2
70to74	16.3
75to79	12.9
80to84	9.7
85+	7.1

**TABLE 32**  
**Life Table For Lancaster White Population**

Age groups	Average Remaining Lifetime
<1	79.1
1to4	78.5
5to9	74.6
10to14	69.7
15to19	64.8
20to24	59.9
25to29	55
30to34	50.1
35to39	45.3
40to44	40.6
45to49	35.9
50to54	31.4
55to59	26.9
60to64	22.5
65to69	18.7
70to74	15.2
75to79	11.9
80to84	9.1
85+	6.8

**TABLE 33**  
**Life Table For Lancaster Black Population**

Age groups	Average Remaining Lifetime
<1	77.3
1to4	78.4
5to9	74.5
10to14	69.5
15to19	64.7
20to24	59.9
25to29	54.9
30to34	50.1
35to39	45.3
40to44	40.6
45to49	36
50to54	31.4
55to59	27.6
60to64	23.1
65to69	19.8
70to74	16.3
75to79	13.8
80to84	10.6
85+	8

**TABLE 34**  
**Life Table For Lancaster American Indian Population**

Age groups	Average Remaining Lifetime
<1	64
1to4	65.5
5to9	61.6
10to14	56.6
15to19	51.6
20to24	47.7
25to29	42.7
30to34	38.1
35to39	33.4
40to44	29.7
45to49	26.4
50to54	22.6
55to59	18.8
60to64	15.8
65to69	14
70to74	10
75to79	6.2
80to84	4.8
85+	3

**TABLE 35**  
**Life Table For Lancaster Asian Population**

Age groups	Average Remaining Lifetime
<1	85.2
1to4	84.8
5to9	80.8
10to14	75.8
15to19	70.8
20to24	65.8
25to29	60.9
30to34	56.2
35to39	51.6
40to44	46.9
45to49	41.9
50to54	36.9
55to59	31.9
60to64	27.3
65to69	22.5
70to74	18.6
75to79	15.5
80to84	11.1
85+	9



**TABLE 36. SUMMARY OF RESIDENT DEATH STATISTICS FROM 1984 TO 2000, LANCASTER COUNTY, NE, 2000**

	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000
<b>NUMBER OF DEATHS</b>																	
Lancaster County	1365	1432	1418	1399	1487	1608	1600	1494	1494	1512	1617	1637	1675	1663	1662	1663	
Lincoln	1243	1331	1302	1274	1387	1394	1371	1370	1339	1410	1483	1409	1558	1438	1468	1520	1528
Balance of County	122	101	116	125	100	114	129	124	155	102	134	128	117	125	114	162	125
<b>RACE</b>																	
White	98.20%	97.40%	97.30%	97.50%	97.90%	97.90%	97.90%	98.20%	97.60%	97.40%	97.80%	97.70%	96.20%	96.20%	97.70%	96.70%	97.2%
Black	1.40%	1.70%	1.60%	1.20%	1.30%	1.50%	1.70%	1.30%	1.60%	1.80%	1.50%	1.40%	2.40%	2.20%	1.30%	1.40%	1.5%
Other	0.40%	0.80%	0.80%	1.30%	0.80%	1.00%	0.40%	0.50%	0.80%	0.90%	0.70%	0.90%	1.40%	1.50%	1.00%	1.80%	1.2%
<b>SEX</b>																	
Male	48.80%	50.00%	49.50%	48.90%	50.40%	50.10%	47.80%	46.50%	49.60%	48.30%	47.80%	47.50%	48.40%	46.40%	47.60%	46.20%	47.1%
Female	51.20%	50.00%	50.50%	51.10%	49.60%	49.90%	52.20%	54.50%	50.40%	51.70%	52.20%	52.50%	51.60%	53.60%	52.40%	53.80%	52.9%
<b>AGE</b>																	
Under 1	2.30%	2.20%	2.10%	2.70%	2.10%	1.70%	1.50%	1.70%	1.60%	2.10%	1.40%	1.30%	1.60%	1.20%	1.60%	1.00%	1.33%
1-14	0.70%	0.60%	0.80%	1.30%	0.40%	0.50%	1.10%	0.80%	0.90%	0.50%	0.60%	0.30%	0.70%	0.40%	0.80%	1.30%	0.54%
15-29	2.40%	2.80%	3.10%	2.40%	2.20%	3.20%	2.50%	1.20%	2.80%	2.40%	2.00%	2.10%	2.10%	1.80%	1.50%	2.00%	1.69%
30-44	2.80%	2.90%	3.70%	2.90%	3.90%	3.10%	4.50%	4.50%	5.20%	4.80%	6.10%	4.50%	4.40%	5.50%	4.40%	4.80%	4.29%
45-59	9.60%	8.30%	9.30%	8.40%	9.10%	8.40%	8.40%	7.90%	7.90%	9.50%	8.80%	7.30%	9.60%	9.70%	10.30%	8.10%	9.01%
60 & Over	82.10%	83.00%	81.00%	82.30%	82.30%	83.00%	81.90%	83.90%	81.60%	80.80%	80.80%	84.40%	81.60%	81.40%	81.40%	82.80%	83.12%
<b>AVERAGE AGE AT DEATH</b>																	
Male	67.4	68.3	67.2	67.9	68	69	67.6	69	68.2	67.8	67.9	69.5	68.3	69.2	68.8	68.8	68.7
Female	76.3	75.2	75	75.4	76	75.1	76.3	77.2	75.1	76.4	76.1	77.9	77.6	77.1	76.4	76.6	77.8
Total	71.9	71.7	71.2	71.8	72	72	72.1	73.5	71.7	72.2	72.1	73.9	73.1	73.4	72.8	73.3	73.9
<b>DEATH RATE</b>																	
Deaths per 1000 Population	6.7	7.1	7	6.7	7	7	7	6.9	6.9	6.9	7.3	6.7	7.2	6.7	6.7	7.07	6.6
INFANT DEATHS	32	32	30	38	31	26	23	25	24	31	22	20	27	18	26	23	22
INFANT MORTALITY RATE (Deaths per 1000 Live Births)	10	10.3	10	12.6	10.4	8.2	7.4	7.8	7.8	9.6	7	6.3	8.2	5.6	7.7	6.4	5.9
NEONATAL DEATHS	19	23	22	22	16	16	13	13	10	19	16	17	17	13	21	16	11
NEONATAL DEATH RATE	5.9	7.4	7.4	6	5.4	5	7.4	4.1	3.3	5.9	5.1	5.4	5.2	4	6.2	4.4	2.9
FETAL DEATHS	25	27	23	19	26	14	23	19	18	23	23	17	20	18	15	21	16
FETAL DEATH RATE	7.8	8.7	7.7	6.3	8.8	4.4	7.4	6	5.9	7.1	7.4	5.4	6.1	5.6	4.4	5.8	4.3

*Lincoln-Lancaster County Health Department*

